

Evaluation of fungicides for management of whit mold of tomato, fall 2011.

Treatment, rate/A (application)^z	Stem lesion length (cm)
Switch, 14 oz (Foliar spray; 3).....	10.3 bc ^y
Inspire Super, 20 fl oz (Foliar spray; 3).....	11.0 bc
Endura, 7 oz (Drip; 3).....	11.5 bc
Endura, 10.5 oz (Drip; 3).....	10.6 bc
Endura, 7 oz (Foliar spray; 3).....	11.0 bc
Endura, 10.5 oz (Foliar spray; 3).....	9.88 bc
Luna Privilege, 6.84 oz (Drip; 3).....	12.2 b
Luna Privilege, 6.84 oz (Drip; 2).....	10.3 bc
Luna Privilege, 6.84 oz (Drip; 1).....	8.66 c
Fontelis, 24 fl oz (Foliar spray; 3)	11.9 b
Heads-up, 0.13 oz/gal (Foliar spray; 3).....	11.9 b
Non-treated control.....	15.6 a
<i>P</i> > <i>F</i>	0.023

^z Listed treatments were applied over a 3 week period at per acre rate basis unless noted otherwise. Stems were inoculated 1 day after last application date, and lesion lengths were evaluated approximately 2 weeks later.

^y Values followed by the same letter are not statistically significant ($P = 0.05$) according to Fisher's LSD test.