RESTRICTED USE PESTICIDE

Due to acute toxicity to humans, aquatic organisms, and avian species. For retail sale to and use only by certified applicators or persons under their direct supervision, and only for those uses covered by the certified applicator's certification.

SUPPLEMENTALLABELING



Insecticide

(EPA Reg. No. 19713-399)

FOR USE IN COMMERCIALLY GROWN ORNAMENTALS (GROWN OUTDOORS IN NURSERIES ONLY)

READ THE ENTIRE DREXELENDOSULFAN 3EC LABEL BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABEL

KEEP OUT OF REACH OF CHILDREN





Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. This labeling and the EPA registered labeling must be in the possession of the user at time of pesticide application.

All applicable directions, restrictions and use precautions on Drexel ENDOSULFAN 3EC LABEL(s) are to be followed. This labeling contains supplemental instructions and restrictions for use of this product which may not appear on the Drexel ENDOSULFAN 3EC label. Follow the instructions carefully.

Crop	Pest Controlled	Rate of Application
Leatherleaf fern (Leather Holly fern) (Except California)	Leatherleaf Fern Borer	1.33 pints per 100 gals. of water (1.3 teaspoons per gal. of water.)
	SPECIFIC DIRECTIONS: Begin treatment when first larval feeding is observed in the mid-vein area at the base of the leaflets. Repeat at 2 to 3 week intervals as necessary.	

USE PRECAUTIONS AND RESTRICTIONS: Food utensils such as teaspoons must not be used for food purposes after use in measuring pesticides. Do not exceed a maximum of 3 pounds of active ingredient (8 pints of this product) per acre per year. Do not use in greenhouses. Do not apply this product through any type of irrigation system.

